

(19)



Eur. patentes Patentamt
Europ. an Patent Office
Office européen des brevets



(11)

EP 1 044 690 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
18.10.2000 Bulletin 2000/42

(51) Int. Cl.: A61K 39/395, C07K 16/06

(21) Application number: 00500066.6

(22) Date of filing: 12.04.2000

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

- Rodríguez Cansel, Carmen
08016 Barcelona (ES)
- Rodenas Navarro, Jesus
08031 Barcelona (ES)
- Saborido Martín, María de las Nieves
08470 San Celoni (Barcelona) (ES)

(30) Priority: 13.04.1999 ES 8900767
11.04.2000 ES 200000533

- (74) Representative: Currell Agullà, Marcelino et al
Dr. Ing. M. Currell Sunol I. S.L.
Agentes de la Propiedad Industrial
Passeig de Gràcia, 66 bis
E-08008 Barcelona (ES)

(71) Applicant: APC Europe S.A.
08016 Barcelona (ES)

(72) Inventors:
• Polo Pozo, Francisco Javier
08210 Barbera del Valles (Barcelona) (ES)

(54) Applications of gamma globulin-rich plasma protein mixtures of animal origin and process for the preparation thereof

(57) The invention concerns applications of mixtures of gamma globulin-rich plasma proteins of animal origin for the preparation of medicinal compositions, food and dietary preparations: [a] suitable for the prevention and treatment of gastrointestinal disorders in human beings and animals; [b] suitable for improving the immunological state of human beings and animals; [c] suitable for the treatment of malabsorption processes in human beings and animals, and [d] suitable for the treatment of clinical conditions accompanied by malnutrition in human beings and animals and a process for

the preparation of the above mixtures comprising diluting a first gamma globulin concentrate in water at a ratio of 1:7 - 1:11; adjusting the pH to 3-6; adding polyethylene glycol at a rate of 2-6 wt%; centrifuging at an acceleration of 3000-11000 g to give a first clarified gamma globulin-rich fraction, which is separated; adjusting the pH thereof to 6-9, adding of a further amount of polyethylene glycol at a rate of 6-12 wt%; centrifuging for a second time at an acceleration of 3000-11000 g to give a second insolubilized fraction, comprising mainly gamma globulin and a second clarified fraction, which are separated.

Description

Field of the Invention

- 5 [0001] This invention relates to applications of a mixture of plasma proteins of animal origin having a gamma globulin content of over 25 wt%, for the preparation of medicinal compositions, food and dietary preparations. It also relates to a process for the preparation of a mixture of plasma proteins of animal origin having a gamma globulin content of over 90 wt%, without the use of chromatographic columns, from a first concentrate, the gamma globulin/proteins ratio of which is over 90 wt%, said first concentrate being derived from animal plasma which has been subjected to at least
- 10 one prior centrifugation operation.

Prior Art Reference

- [0002] On the basis of the known literature, the situation relating to the fractionation process is understood to be as follows:
- 15 [0003] In "A chromatographic Procedure for the Purification of Human Plasma Albumin", Curling, J.M., J. Berglöf, L. O. Lindquist & S. Eriksson; Vox Sang. 33:97-107 (1977), there is described a fractionation method starting out from human plasma, in which polyethylene glycol (PEG) 4000 is used to obtain a gamma globulin-rich precipitate and an albumin fraction concentrate, which is finally purified to 99% purity using a chromatographic system.
- 20 [0004] In "A Heat Polyethylene Glycol Procedure as an Alternative Method for Plasmafractionation", Guddat, W. & K. Hülger; Folia Haematol., Leipzig 108 (1982) 6, S. 840-855, there is described a fractionation process starting out from human plasma for preparing albumin of high purity. In the process described, fractionation steps with PEG 4000 are alternated with the use of sodium octanoate, with heating to high temperatures (75°C) to obtain the albumin-enriched fraction.
- 25 [0005] In "A new high quality Albumin for therapeutic use", Joint WHO/IAS Symposium on the Standardization of Albumin, Plasma Substitutes and Plasmapheresis, Hansen, J.F. & M. Erban. Geneva 1980, Develop. Biol. Standard. 48, pp. 105-112, there is described a fractionation method for the preparation of human albumin for therapeutic use. In 48, pp. 105-112, there is described a fractionation method for the preparation of human albumin for therapeutic use. In human beings, by way of injection, in which combined fractionation steps with PEG 3000 and the use of caprylic acid, temperature and ethanol are detailed, for the preparation of different plasma and high purity albumin fractions.
- 30 [0006] In "Fractional Precipitation of Proteins with Polyethylene Glycol", Hao, Y.L., K.C. Ingham & M. Wickehauser, there is cited a fractionation process on human plasma for preparing different plasma fractions. According to the process described, PEG 4000 is used for a first separation, followed by purification steps in chromatographic columns.
- 35 [0007] In "Optimization for Selective Fractionation of Bovine Blood Plasma Proteins Using Poly(ethylene glycol)", Lee, Y.Z., T. Aishima, S. Nakai & J.S. Sim J. Agri. Food Chem. 1987, 35, 958-962, there is described a plasma fractionation process carried out on bovine plasma. By working with different PEG 4000 and sodium chloride concentrations and at certain pH values, it was possible to separate three major fractions: fibrinogen, immunoglobulins and albumin.
- 40 [0008] In "Fractionation of Plasma with Polyethylene Glycol", Polson, A. & Ruiz-Bravo; Vox Sang. 23, 107-118 (1972) there is described a fractionation process on human plasma using PEG 6000 for preparing different protein fractions included in the plasma. According to these workers, the process may be carried out working with PEG alone or in combination with organic salts or agents, or using chromatographic membranes.
- 45 [0009] In "An Alternative Method of Large Scale Plasma Fractionation for the Isolation of Serum Albumin", Schneider, W., H. Lefèvre, H. Fiedler & L.J. McCarty; Blut, Band 30, Seite 121-134 (1975), there is mentioned the possibility of obtaining high purity albumin from human plasma using sodium octanoate in combination with ethanol and high temperatures (68°C) for the separation of albumin from the remaining plasma proteins. Subsequently, the albumin is precipitated with the use of PEG 4000 and after resuspension in water a liquid form is obtained, or a powder is obtained after freeze-drying.
- 50 [0010] In "The Isolation of IgG from Mammalian Sera with the Aid of Caprylic Acid", Steinbuch, M. & R. Audren; Archives of Biochemistry and Biophysics 134, 279-284, (1969), there is related a fractionation process on human, ovine, goat or rabbit plasma using caprylic acid (octanoic acid) for purification of the immunoglobulin-rich fraction. After desalination by dialysis, the fraction was subsequently concentrated or obtained in powder form through a freeze-drying process.
- 55 [0011] In "Precipitation of Plasma Lipoproteins by PEG-6000 and Its Evaluation with Electrophoresis and Ultracentrifugation", Vilkarl, J.; Scand. J. Clin. Lab. Invest., Vol. 38, 1976, there is related a fractionation method on human plasma using different levels of PEG 6000 for the preparation of the lipoprotein-rich (LDL, VLDL, HDL) fractions in a single precipitation step.
- [0012] Finally, in "Large Scale Preparation of Macroglobulins", Wickehauser, M. and Y.L. Hao; Vox Sang. 23: 119-125 (1972), there is mentioned the possibility of fractionation of an immunoglobulin M-rich (IgM) fraction from the fraction prepared from human plasma, using Conn's aqueous alcoholic fractionating method. In this case, starting out from

the precipitate III of Cohn's method and using PEG 6000 together with zinc sulphate, they obtained a supernatant which they subsequently precipitated with ammonium sulphate.

the precipitate III of Conn's method and being the only one which they subsequently precipitated with ammonium sulphate. Conn [19013] The literature covering the period from the late 80's up to date, apart from being very scarce, does not provide any more information on this subject.

[0013] The literature covering the plasma components expressed in the foregoing publications significantly different concepts from the papers cited, there is disclosed a process in which, starting out from an animal

[0014] According to some of the papers cited and subjecting it to a first prior centrifugation, under certain conditions, there is prepared an insolubilized nitrogen-rich fraction and a clarified fraction containing the remaining proteins contained in the plasma and which are water soluble. After a certain treatment, this clarified fraction is subjected to a second prior centrifugation, from which there is separated a new concentrated gamma globulin-rich fraction and a new clarified

albumin-rich fraction.

of these various protein mixtures in human beings are known, for example in

[0015] Certain applications of fibrinogen-rich plasma protein mixtures in human beings are known, for example in the treatment of haemophiliacs. Nevertheless, these applications require intravenous application, which means that the plasma must be transfused to the patient, to avoid incompatibility problems.

the treatment of haemophiliacs, the origin of the mixture has to be human, to avoid incompatibility problems. [0016] Also known are certain applications of albumin-rich plasma protein mixtures, for example in surgery. Nevertheless, the application is also intravenous and the origin of the mixture has to be human.

Summary of the Invention

[0017] It is an object of the invention to provide applications of a mixture of plasma proteins of animal origin having a gamma globulin content of over 25 wt%, for the preparation of medicinal compositions, food and dietary preparations; a gamma globulin content of over 25 wt%, for the preparation of medicinal compositions in human beings and animals; [b] suitable for [a] suitable for the prevention and treatment of gastrointestinal disorders in human beings and animals; [c] suitable for improving the immunological state of human beings and animals; particularly the small intestine syndrome; refractory diarrhoea of autoimmune origin; lymphoma; post-gastroctomy steatorrhea; carcinoma of the pancreas; wide pancreatic resection; mesenteric vascular insufficiency; amyloidosis; scleroderma; eosinophilic enteritis; and [d] suitable for the treatment of clinical conditions accompanied by malnutrition in human beings and animals, particularly ulcerative colitis; Crohn's disease; cancerous cachexia for chronic enteritis caused by chemo- and/or radiotherapy; infectious medical pathology involving severe malabsorption; AIDS; cystic fibrosis; low flow enterocutaneous fistulas; and Infantile renal insufficiency endangering patient growth.

[0018] The plasma proteins are preferably obtained from porcine, bovine, goat, equine and ovine plasma. The use of these plasma proteins is preferred, above all for the treatment of human beings.

[0018] The plasma proteins are particularly preferred, above all for the treatment of human beings.
[0019] Of porcine plasma is particularly preferred, above all for the treatment of human beings.
[0020] One advantageous form of application is obtained when the medicinal compositions, food and dietary preparations are usable dissolved in water, milk, juices, yoghurts and other foodstuffs. These other foodstuffs are preferably

[0020] Preferably any variety of fibres, prebiotic, probiotic products, vitamins and minerals are added to the medicinal compositions, food and dietary preparations.

[0021] It is of particular interest that the immuno-globulins should be non-specific, since this involves a large cost reduction, they have a therapeutic activity sufficient for slight infections and are an aid in more serious infections and to mask any of the symptoms of the infection to be treated, whereby they may be used immediately such symptoms

appear. ¹ This effect is obtained when their use is combined with the administration of anti-

[0022] Finally, a further preferred application is obtained when their use is combined with the use of a gamma globulin fraction of over 90 wt%.

[0023] It is also an object of the invention to prepare a concentrate of the gentamicin solution, since in such case smaller amounts are required for the later applications thereof, by way of a process significantly simpler and less expensive in comparison with the regular chromatographic techniques. This object is achieved with the aid of a process according to the invention, which is described in detail in the following.

reducing the production cost in comparison with the regular chromatographic technique. The process of the type first mentioned above, characterized in that: [i] said first concentrate is diluted in water in a first process: water ratio ranging from 1.7 - 1.11, giving a first mixture; [ii] the pH of said first mixture is adjusted for a second process: water ratio ranging from 3 - 6, to give a second mixture; [iii] from 2 to 6 % of polyethylene glycol relative to the first time to a value ranging from 3 - 6, to give a second mixture; [iv] said third mixture is centrifuged for a first time at an acceleration ranging from 3000 to 11000 g, giving a first insolubilized beta globulin-rich fraction and a first clarified gamma globulin-rich fraction, both first fractions being separated; [v] the pH of said gamma globulin-rich fraction is adjusted for a second time to a value ranging from 6-12 w/v relative to said fourth mixture, to which polyethylene glycol is added to a weight proportion ranging from 6-12 w/v relative to said fourth mixture, to provide a fifth mixture; [vi] said fifth mixture is centrifuged for a second time at an acceleration ranging from 3000 - 11000 g to give a second insolubilized fraction composed mainly of gamma globulin and a second clarified fraction, both second fractions being separated.

[0024] The animal plasma used in this process is preferably porcine, bovine, ovine, equine or goat plasma, with the porcine plasma being particularly preferred.

(0025) Concentrates of the product to which the invention relates may be obtained with the use of spray drying driers.

freeze-drying or any other drying technique. The present invention does not exclude the preparation of these concentrates in liquid or frozen form. They may also be prepared by microencapsulation of the active ingredient, protecting it and stabilizing it against high temperatures, oxidants, pH, light, humidity, etc.

[0026] They may be supplied in tablet, capsule, drinkable vial, granulated, powder, cream form, both as a sole ingredient and in association with other excipients or active ingredients.

Detailed Description of the Invention

[0027] The invention consists of the application of immuno-globulin concentrates, particularly of porcine origin, for the prevention and co-treatment of gastrointestinal disorders of diverse origin. These concentrates are preferably of non-specific immuno-globulins. It is contemplated that they are administered orally or topically and are used both alone and in association with other substances.

[0028] The use of porcine immuno-globulins reduces the risk of allergy reactions caused by other immuno-globulin sources, having an allergenic risk. This allergenic risk appears, for example, due to the fact that both cow's milk and egg are foodstuffs used in infant nutrition in the first stages of replacement of the breast milk. This means that certain individuals have developed allergies specific to proteins from these sources.

[0029] The majority of processes in which specific immuno-globulins are used are costly, since animals must be hyperimmunized with specific antigens. These treatments require a precise prior diagnosis, which is costly both financially and in the time required for obtaining the results of the analyses.

[0030] The application of non-specific immuno-globulins (which contain antibodies for a very high number of micro-organisms) is advantageous in light intestinal infections where the presence of antibodies available in the immuno-globulin concentrate is already sufficient to stop the symptoms caused by the infection. In this way, the symptoms may also be alleviated of more serious infections, in which it is necessary to have recourse to treatment with specific antibiotics after diagnosis, since they do not mask any of the symptoms of the infection and, therefore, may be applied immediately after the appearance of said symptoms.

[0031] The administration of non-specific immuno-globulins of porcine origin orally or enterally is particularly advantageous, since it does not cause immunological rejection problems since the said immuno-globulins do not pass through the intestinal wall and since they perform their protective or therapeutic activity locally in the intestinal passage.

[0032] It frequently happens that, in population groups having a risk of gastrointestinal disorders, such as a new born baby of low weight at birth, new born babies not breast-feeding, elderly people with immuno deficiencies, adults having an impaired immunity system, particularly in the case of AIDS, persons undergoing chemotherapy in cancer processes, or persons who have undergone a bone medulla transplant, gastrointestinal infections are chronic and are often accompanied by opportunistic infections taking advantage of the imbalance of the intestinal flora to manifest their pathogeny. In such cases, it usually happens that the therapies used are not completely effective, since in the best of cases they manage to reduce the effect caused by the first pathogen, but are inactive against the opportunistic germs, whereby the use of non-specific gamma globulin concentrates of broad antibacterial spectrum is very advisable in these situations, since further to helping to restore the bacterial ecosystem of the intestine, it prevents the action of opportunistic pathogens and thereby helps in the recovery of the intestinal epithelium. It is for this same reason that the joint application of an immuno-globulin enriched preparation with the specific therapy is of interest.

[0033] The application of porcine immuno-globulin enriched preparations affords the additional advantage that it does not cause the appearance of a secondary symptomatology associated with the use of other compounds which are used in processes of gastrointestinal disorders, such as are constipation, loss of gastrointestinal flora or reduction of appetite, since, with their being natural proteins, after their effect associated with the bioactivity of such molecules, they will be degraded or excreted in the same way as happens with the remaining animal proteins ingested daily in the diet.

[0034] Since it is a case of a high spectrum immuno-globulin concentrates, obtained from a very large volume of blood of animals from different sources, it is to be expected that it will contain antibodies for a very varied number of bacteria and virus. In human medicine, the treatment of viral gastrointestinal infections has not yet been solved, since the antibiotics do not have an effect on these virus, whereby the use of immuno-globulins could mean an advance in the reduction of the acuity of such infections.

[0035] It is well known that both bacteria and virus have the capacity of becoming resistant to antibiotics or chemical preparations. Mammals combat such resistance by creating new antibodies specific to the new mutation of the bacteria. In such a way that the bacterial or viral mutations are naturally corresponded to by the appearance of antibodies or predominant in the blood against such mutations. In these cases, it usually happens that with time the antibodies or immuno-globulins obtained from hyperimmunized animals cease to have a complete effect and new generations of antibiotics or specific immuno-globulins have to be obtained. This does not happen in our case, since with the immuno-globulin enriched preparation being obtained from sacrificed animals, it contains antibodies against the new strains of bacteria that have mutated to acquire resistance to the antibiotics. In this way, they have at all times an immunological

activity against the subclasses of pathogens which cause the gastrointestinal disorders.

[0036] These applications are also appropriate for pets. The microbiological quality of their food is lower than that of human food and above all is due to the fact that the food usually remains in the feeder for many hours at room temperature, whereby bacterial growth may develop under such conditions, and may cause intestinal infections. The use of specific immuno-globulins from hyperimmunized animals for the treatment of these gastrointestinal infections is infrequent because of the high cost involved in such therapy, as well as the lack of a precise diagnosis of such infection. Therefore, the use of broad bacterial and viral spectrum immuno-globulin concentrates, particularly those of porcine origin, is an economical and viable alternative for use in reducing the seriousness of gastrointestinal infections and recovering the animals' intestinal balance, as well as a preventive use in animals having chronic infections and, therefore, poor intestinal absorption.

[0037] It is a regular practice to include powdered plasma in the initiation and weaning diets of farm animals, particularly pigs. This affords benefits with regard to growth and reduction of diarrhoeas and mortality in this period of life. Such benefits have been associated in the scientific literature with the immuno-globulin fraction present in the powdered plasma, whereby the use of an immuno-globulin-rich fraction in the feeds administered at these ages will have a higher effect to that caused by the whole plasma. Pigs belong to a group of mammals born immunodeficient which need to ingest the maternal colostrum, rich in immuno-globulins, to incorporate passive immunity into their organism. This must occur during the first 24-36 hours of life, which is when there is an intestinal permeability to macromolecules, whereby part of the immuno-globulins ingested in the colostrum will pass into the sucking animal's blood, providing it with passive immunity for the first weeks, until the animal is capable of generating its own immunitary system.

[0038] Nevertheless, the supplies of immuno-globulins as substitutes for the maternal colostrum known on the market are based on bovine immuno-globulins. The fact of using porcine origin immuno-globulins represents an advantage over the use of bovine immuno-globulins, since the action time in blood of bovine immuno-globulins is substantially inferior to that of porcine immuno-globulins, whereby the passive immunity of the sucking animals is improved during the first 24 hours of life.

[0039] In some applications, it is convenient to have plasma protein mixtures containing gamma globulin concentrations of over 90%.

[0040] Nevertheless, in other applications, it is sufficient to have plasma protein mixtures containing gamma globulin concentrations, relative to total proteins, of over 25%. In these cases, said mixtures may be prepared in a cheaper way and, additionally, these mixtures retain other proteins, peptides, vitamins and growth factors of high value and which are removed in the plasma protein mixtures whose gamma globulin content is over 90%.

[0041] For a better understanding of the invention, there are disclosed hereinafter examples of the diverse steps of the process of the invention, with inclusion of steps prior to the preparation of the starting material.

Example 1

Separation of first fractions from animal plasma

[0042] 1000 kg of unhaemolized porcine plasma were charged in a work reactor at 4°C and pH 7.7. Stirring was applied at 50 r.p.m. and 53.18 kg of hydrochloric acid were added, whereby the pH was lowered to 4.5. 55 kg of polyethylene glycol (PEG) 4000 were added, whereby the mixture contained 6 wt% of PEG. The pH was readjusted to 4.5. The reaction mixture was left under stirring for 5 hours and thereafter was centrifuged in a centrifuge at 9000 g, with a bowl for recovering the concentrated fraction therein. After centrifugation, 33.7 kg were obtained of a fibrinogen-rich fraction and 1052 kg of a clarified fraction containing the remaining soluble proteins.

[0043] After removal of the polyethylene glycol, the clarified fraction had a gamma globulin concentration, relative to total proteins, of approximately 30%.

Example 2

Treatment of the clarified fraction obtained

[0044] The 1052 kg of clarified fraction prepared according to Example 1 were held under constant stirring at 4°C and the pH was adjusted to 7.0 by addition of 33.32 kg of sodium hydroxide. Thereafter, 89.2 kg of PEG were added to maintain the wt% at 12, the pH being adjusted once again to 7.0. The mixture was left under constant stirring at the temperature indicated for 5 hours. Thereafter, the mixture was centrifuged in the same centrifuge as mentioned in the previous example and 1093 kg of a new clarified fraction and 54.8 kg of a gamma globulin-rich concentrate were obtained.

[0045] This gamma globulin-rich concentrate had a gamma globulin concentration relative to total proteins of approximately 50%.

Example 3Treatment of the gamma globulin-rich concentrate

[0046] To the 54.8 kg of gamma globulin-rich concentrate obtained according to the previous example were added 492 kg of water (concentrate:water ratio \rightarrow 1:9). The pH was adjusted to 8.0 by addition of 7.7 kg of hydrochloric acid and a further 32.8 kg of PEG were added to keep the PEG concentration at 6 wt%, the pH being adjusted again with hydrochloric acid. The mixture was held under constant stirring at 4°C for 5 hours, after which it was centrifuged in a centrifuge of the type indicated in the previous examples. After the centrifugation process, 17 kg were obtained of a beta globulin-rich insolubilized fraction and 552 kg of a clarified gamma globulin-rich fraction.

Example 4Treatment of the new clarified fraction obtained

[0047] To the 552 kg of the clarified gamma globulin-rich fraction obtained according to the previous example there were added 4.8 kg of sodium hydroxide, whereby the pH thereof was adjusted to 9.0. Furthermore, 36 kg of PEG were added, whereby the proportion of PEG (bearing in mind the PEG contained in the fraction) was held at 12 wt%. The pH was adjusted again to 8.0. The mixture was held under constant stirring at 4°C for 5 hours, after which it was centrifuged in an apparatus of the type mentioned above and thus 550 kg of a clarified fraction and 30 kg of a fraction having a high gamma globulin content were obtained.

Example 5Treatment of the high gamma globulin content fraction

[0048] The 30 kg of the high gamma globulin content fraction obtained according to the previous example were dissolved in water by addition of 60 kg of water (ratio 1:2 by weight). The pH was adjusted to 6.0. Once the mixture was perfectly dissolved, it was dried in an atomizer (spray dried), by removing the water contained in the liquid product by subjecting it to transportation with high temperature air (150°C at the inlet and 100°C at the outlet) for a short period of time, whereby the water was evaporated without affecting the solubility of the protein to be dried. The powdered product was finally packaged. The characteristics of the powdered product obtained were:

Humidity	< 10 %
Protein	90 \pm 2 %
Ashes	< 2 %
Insolubles	< 2 %
Standard proteinogram % (relative to protein)	
Albumin	1.0 %
Alpha globulin	5.3 %
Beta globulin	~ 4.0 %
Gamma globulin	90.63 %

[0049] The utility of the compositions to which reference is made may be due in part to the fact that the product prepared according to the process of the invention may contain IgM, transferrin, cellular growth factors or alpha 2 macroglobulin.

Claim

1. Applications of mixtures of gamma globulin-rich plasma proteins of animal origin having a gamma globulin content of over 25 wt%, for the preparation of medicinal compositions, food and dietary preparations suitable for the prevention and treatment of gastrointestinal disorders in human beings and animals.
2. Applications of mixtures of gamma globulin-rich plasma proteins of animal origin having a gamma globulin content of over 25 wt%, for the preparation of medicinal compositions, food and dietary preparations suitable for improving the immunological state in human beings and animals.
3. Applications of mixtures of gamma globulin-rich plasma proteins of animal origin having a gamma globulin content of over 25 wt%, for the preparation of medicinal compositions, food and dietary preparations suitable for the treatment of malabsorption processes in human beings and animals.
4. The application according to claim 3, characterized in that said malabsorption processes are: small intestine syndrome; refractory diarrhoea of autoimmune origin; lymphoma; post-gastrectomy steatorrhea; carcinoma of the pancreas; wide pancreatic resection; mesenteric vascular insufficiency; amyloidosis; scleroderma; eosinophilic enteritis.
5. Applications of mixtures of gamma globulin-rich plasma proteins of animal origin having a gamma globulin content of over 25 wt%, for the preparation of medicinal compositions, food and dietary preparations suitable for the treatment of clinical conditions with malnutrition in human beings and animals.
6. The application according to claim 5, characterized in that said clinical conditions are: ulcerative colitis; Crohn's disease; cancerous cachexia for chronic enteritis caused by chemo- and/or radiotherapy; infectious medical pathology involving severe malabsorption; AIDS; cystic fibrosis; low flow enterocutaneous fistulas; and infantile renal insufficiency endangering patient growth.
7. The application according to at least one of claims 1 to 6, characterized in that said plasma proteins are obtained from porcine, bovine, goat, equine or ovine plasma.
8. The application according to at least one of claims 1 to 7, characterized in that said medicinal compositions, food and dietary preparations may be used dissolved in water, milk, juices, yoghurt or other foodstuffs.
9. The application according to claim 8, characterized in that said other foodstuffs are humanised milk, paps or other baby foods.
10. The application according to at least one of claims 1 to 9, characterized in that to said medicinal compositions, food and dietary preparations there may be added any variety of fibres, prebiotic, probiotic products, vitamins or minerals.
11. The application according to at least one of claims 1 to 10, characterized in that said medicinal compositions, food and dietary preparations are preferably used with the administration of antibiotics.
12. The application according to at least one of claims 1 to 11, characterized in that said mixtures of plasma proteins are of porcine origin and in that they are applied for the treatment of human beings.
13. The application according to at least one of claims 1 to 12, characterized in that said gamma globulin content is over 90 wt%.
14. A process for the preparation of mixtures of plasma proteins of animal origin having a gamma globulin content of over 90 wt%, without the use of chromatographic columns, from a first concentrate, the gamma globulin/proteins ratio of which is over 30 wt%, said first concentrate being derived from animal plasma which has been subjected to at least one prior centrifugation operation, characterized in that (i) said first concentrate is diluted in water in a first concentrate: water ratio ranging from 1:7 - 1:11, giving a first mixture; (ii) the pH of said first mixture is adjusted for the first time to a value ranging from 3 - 6, to give a second mixture; (iii) from 2 to 8 wt% of polyethylene glycol relative to said second mixture is added thereto, giving a third mixture; (iv) said third mixture is centrifuged for a first time at an acceleration ranging from 3000 to 11000 g, giving a first insolubilized beta globulin-rich fraction and

a first clarified gamma globulin-rich fraction, both first fractions being separated; [v] the pH of said gamma globulin-rich fraction is adjusted for a second time to a value ranging from 6 to 9 to give a fourth mixture, to which polyethylene glycol is added to a weight proportion ranging from 6-12 wt% relative to said fourth mixture, to provide a fifth mixture; [vi] said fifth mixture is centrifuged for a second time at an acceleration ranging from 3000 - 11000 g to give a second insolubilized fraction composed mainly of gamma globulin and a second clarified fraction, both second fractions being separated.

15. The process according to claim 14, characterized in that said animal plasma is porcine, bovine, ovine, equine or goat plasma.
16. The process according to claim 15, characterized in that said animal plasma is porcine plasma.
17. The process according to at least one of claims 14 to 16, characterized in that said second insolubilized fraction is dissolved in water at a second insolubilized fraction/water ratio ranging from 1:2 - 1:9, the pH is adjusted for a third time to a value ranging from 5 to 9 and the fraction is dried.
18. The process according to claim 17, characterized in that said drying is effected in an atomizer, said second insolubilized fraction being subjected to a high temperature air flow.
19. The process according to claim 18, characterized in that said high temperature varies over a range of 150-250°C at the inlet and a range of 70-100°C at the outlet.
20. The process according to at least one of claims 14 to 19, characterized in that said gamma globulin concentrates are suitable for preparation by a microencapsulation process.
21. The process according to at least one of claims 14 to 17, characterized in that said gamma globulin concentrates are prepared in liquid, frozen, concentrated or dried form by a process other than the one using an atomizer.
22. The process according to at least one of claims 14 to 21, characterized in that said first adjustment is effected with hydrochloric or phosphoric acid.
23. The process according to at least one of claims 14 to 22, characterized in that said separation is effected in a filter press with a filtering pore sieve.
24. The process according to at least one of claims 14 to 20, characterized in that said separation is effected in a centrifuge having an acceleration of over 3000 g.
25. The process according to claim 24, characterized in that said centrifuge is continuous.
26. The process according to claim 24, characterized in that said centrifuge has an internal bowl.
27. The process according to at least one of claims 14 to 24, characterized in that said second adjustment is effected with sodium hydroxide.

EP 1 044 690 A1

European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 50 0065

DOCUMENTS CONSIDERED TO BE RELEVANT

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	EP 0 064 210 A (CUTLER LAB) 10 November 1982 (1982-11-10)	1-3, 5, 6	A61K39/395 C07K16/06
Y	* see claims 1 and 7, pages 6-8 and page 10 lines 17-27 *	1-13	
X	US 5 531 989 A (PAUL STEPHEN M) 2 July 1996 (1996-07-02)	1-3, 5-7, 10	A61K
Y	* see abstract and col. 3-4 *	1-13	
X	HASKO F. ET AL.: "Fractionation of plasma proteins with PEG." HAEMATOLOGICA, vol. 14, no. 2, 1981, pages 199-205, XP000917117 Budapest	14, 20, 22, 27	A61K
Y	* see abstract and fig. 2 *	14-27	
Y	PATENT ABSTRACTS OF JAPAN vol. 013, no. 169 (C-587), 21 April 1989 (1989-04-21) & JP 64 000028 A (CHEMO SERO THERAPEUT RES INST), 5 January 1989 (1989-01-05) * abstract *	1-13	A61K
X	PATENT ABSTRACTS OF JAPAN vol. 013, no. 556 (C-664), 11 December 1989 (1989-12-11) & JP 01 230533 A (GREEN CROSS CORP:THE), 14 September 1989 (1989-09-14) * abstract *	14, 20	
Y		14-27	A61K
Y	EP 0 338 229 A (BIOTEST PHARMA GMBH) 25 October 1989 (1989-10-25) * see abstract 3	1-13	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 14 August 2000	Examiner Gora, V
CATEGORY OF CITED DOCUMENTS			
X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background Q: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons S: member of the same patent family, corresponding documents			

EPO FORM 1501 (01/95) PRECIS

EP 1 044 690 A1

European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 50 0066

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of documents with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (InCL17)
Y	SU 1 754 113 A (MO NII EPIDEMIOLOGII MIKROBIOL ; NITEX PATOLOGII TERAPII AMN SS (SU) 15 August 1992 (1992-08-15) * see abstract *	1-13	
Y	BPI, VFA, BAH, VAP: "Rote Liste 1999" 1999, ECV, AULENDORF XP002144418 * see compounds 750003, 75007-750111 and 75013-75014 *	1-27	
Y	TRUSOVA L. ET AL.: "Fractionation of blood serum proteins by non-ionic water-soluble polymers." YSES. GOS. NAUCHNO-KONTROL'N. INST. VET. PREP., 1980, pages 98-101, XP002144417 * see conclusion *	1-27	
			TECHNICAL FIELDS SEARCHED (InCL17)
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 14 August 2000	Examiner Gore, V
CATEGORY OF CITED DOCUMENTS		T: theory or principle underlying the invention E: earlier patent documents, but published on or after the filing date D: document cited in the application L: documents cited for other reasons X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: unpublished document	

DOCUMENT CONTAINED IN THE APPLICATION

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 00 50 0066

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office (EPO) file on the European Patent Office in no way liable for these particulars which are merely given for the purpose of information.

14-08-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0064210 A	10-11-1982	DE 3272680 D	25-09-1986
		JP 1941158 C	23-06-1995
		JP 6069962 B	07-09-1994
		JP 57185222 A	15-11-1982
		US 4477432 A	16-10-1984
US 5531989 A	02-07-1996	US 5531988 A	02-07-1996
		AU 709155 B	19-08-1999
		AU 4013695 A	23-05-1996
		CA 2203762 A	09-05-1996
		EP 0787006 A	06-08-1997
		WO 9613271 A	09-05-1996
		US 5744134 A	28-04-1998
JP 64000028 A	05-01-1989	JP 2672303 B	05-11-1997
JP 01230533 A	14-09-1989	JP 2613409 B	28-05-1997
EP 0338229 A	25-10-1989	AT 90569 T	15-07-1993
		AU 637045 B	20-05-1993
		AU 3417189 A	24-11-1989
		BR 8907385 A	21-05-1991
		CN 1037276 A	22-11-1989
		DD 283937 A	31-10-1990
		WO 8910139 A	02-11-1989
		DE 58904683 D	22-07-1993
		DK 251990 A	18-10-1990
		IN 168629 A	11-05-1991
		JP 3504718 T	17-10-1991
		NO 904440 A	15-10-1990
		NZ 228595 A	26-11-1991
		PT 90318 A	10-11-1989
		ZA 8902855 A	27-12-1989
SU 1754113 A	15-08-1992	NONE	

EP 00 50 0066

For more details about this annex: see Official Journal of the European Patent Office, No. 12/82